

## ON THE OPTIMAL MULTILINEAR BOHNENBLUST–HILLE CONSTANTS

D. NUÑEZ-ALARCÓN\*, D. PELLEGRINO\*\*, J.B. SEOANE-SEPÚLVEDA\*\*\*,  
AND D. M. SERRANO-RODRÍGUEZ\*

ABSTRACT. The upper estimates for the optimal constants of the multilinear Bohnenblust–Hille inequality obtained in [2] are here improved to:

- (1) For real scalars:  $K_n \leq \sqrt{2} (n-1)^{0.526322}$ .
- (2) For complex scalars:  $K_n \leq \frac{2}{\sqrt{\pi}} (n-1)^{0.304975}$ .

We also obtain sharper estimates for higher values of  $n$ . For instance,

$$K_n < 1.30379 (n-1)^{0.526322}$$

for real scalars and  $n > 2^8$  and

$$K_n < 0.99137 (n-1)^{0.304975}$$

for complex scalars and  $n > 2^{15}$ .

## 1. PRELIMINARIES. FIRST ESTIMATES

Let  $\mathbb{K}$  be the real or complex scalar field. The multilinear Bohnenblust–Hille inequality asserts that for every positive integer  $n \geq 1$  there exists a sequence of positive scalars  $(C_n)_{n=1}^\infty$  in  $[1, \infty)$  such that

$$\left( \sum_{i_1, \dots, i_n=1}^N |U(e_{i_1}, \dots, e_{i_n})|^{\frac{2n}{n+1}} \right)^{\frac{n+1}{2n}} \leq C_n \sup_{z_1, \dots, z_n \in \mathbb{D}^N} |U(z_1, \dots, z_n)|$$

for all  $n$ -linear forms  $U : \mathbb{K}^N \times \dots \times \mathbb{K}^N \rightarrow \mathbb{K}$  and every positive integer  $N$ , where  $(e_i)_{i=1}^N$  denotes the canonical basis of  $\mathbb{K}^N$  and  $\mathbb{D}^N$  represents the open unit polydisc in  $\mathbb{K}^N$ . In the last years a considerable effort was spent in the searching of optimal values for the constants  $C_n$ ; for details and the state-of-the-art we refer to [2]. The notation and terminology used in this note are the same as those from [2], where it is proved that the optimal multilinear Bohnenblust–Hille constants  $(K_n)_{n=2}^\infty$  satisfy

$$(1.1) \quad K_n < 1.65 (n-1)^{0.526322} + 0.13 \text{ (real scalars)}$$

and

$$K_n < 1.41 (n-1)^{0.304975} - 0.04 \text{ (complex scalars)}.$$

The proof of the above estimates is achieved by following a series of technical steps. In the case of real scalars, using some previous lemmata, it is observed that the sequence

$$M_n = \begin{cases} (\sqrt{2})^{n-1} & \text{if } n = 1, 2 \\ DM_{\frac{n}{2}} & \text{if } n \text{ is even, and} \\ DM_{\frac{n+1}{2}} & \text{if } n \text{ is odd} \end{cases}$$

---

\*Supported by Capes.

\*\*Supported by CNPq Grant 301237/2009-3.

\*\*\*Supported by grant MTM2012-34341.

satisfies the multilinear Bohnenblust–Hille inequality, where  $D = \frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}$ . Then, using a “uniform approximation” argument, the estimate (1.1) is achieved. In this section we remark that this final step of the proof, i.e., the uniform approximation argument, can be dropped and a quite simple argument provides even better constants. In fact, from [2] we know that, for all  $k \geq 1$  and  $n \geq 2$ , we have

$$M_n = \sqrt{2}D^{k-1} \text{ whenever } n \in B_k = \{2^{k-1} + 1, \dots, 2^k\}.$$

Thus,  $k - 1 \leq \log_2(n - 1)$  and, hence,

$$M_n \leq \sqrt{2}D^{\log_2(n-1)} = \sqrt{2}(n-1)^{\log_2\left(\frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}\right)} \leq \sqrt{2}(n-1)^{0.526322}.$$

Using a similar argument (for complex scalars) it follows that

$$M_n \leq \frac{2}{\sqrt{\pi}}(n-1)^{\log_2\left(e^{\frac{1}{2}-\frac{1}{2}\gamma}\right)} \leq \frac{2}{\sqrt{\pi}}(n-1)^{0.304975}$$

for the complex scalar field. Of course, the other estimates of [2] related to the above results can be straightforwardly improved by using these new estimates.

## 2. SHARPER ESTIMATES FOR BIG VALUES OF $n$

In this section we improve our previous estimates for *large* values of  $n$ .

**2.1. Real case.** If  $(C_n)_{n=1}^\infty$  denotes the sequence in [2, (4.3)], if we fix any  $k_0$ , it is obvious that

$$J_n = \begin{cases} C_n & \text{if } n \leq 2^{k_0}, \\ DJ_{\frac{n}{2}} & \text{if } n > 2^{k_0} \text{ is even, and} \\ D\left(J_{\frac{n-1}{2}}\right)^{\frac{n-1}{2n}} \left(J_{\frac{n+1}{2}}\right)^{\frac{n+1}{2n}} & \text{if } n > 2^{k_0} \text{ is odd} \end{cases}$$

with  $D = \frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}$ , satisfies the multilinear Bohnenblust–Hille inequality. For  $n > 2^{k_0}$ , let  $k_1 > k_0$  be such that

$$2^{k_1-1} + 1 \leq n \leq 2^{k_1}.$$

Then

$$k_1 - k_0 \leq \log_2\left(\frac{n-1}{2^{k_0-1}}\right)$$

and

$$\begin{aligned} K_n &\leq J_{2^{k_1}} = D^{k_1-k_0} C_{2^{k_0}} \\ &\leq C_{2^{k_0}} D^{\log_2\left(\frac{n-1}{2^{k_0-1}}\right)} \\ &= \frac{C_{2^{k_0}}}{D^{k_0-1}} (n-1)^{\log_2 D}. \end{aligned}$$

We thus have

$$K_n \leq \frac{C_{2^{k_0}}}{D^{k_0-1}} (n-1)^{\log_2\left(\frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}\right)}.$$

From [3, Theorem 3.1] we know that

$$(2.1) \quad C_{2^{k_0}} \leq 4D^{k_0-4}$$

whenever  $k_0 \geq 4$ . Thus,

$$K_n \leq \frac{4}{\left(\frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}\right)^3} (n-1)^{\log_2 \left(\frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}\right)} < 1.338887 (n-1)^{0.526322}.$$

Summarizing, we have:

**Theorem 2.1.** *If  $n > 16$ , then*

$$K_n \leq \frac{4}{\left(\frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}\right)^3} (n-1)^{\log_2 \left(\frac{e^{1-\frac{1}{2}\gamma}}{\sqrt{2}}\right)}.$$

Numerically,

$$(2.2) \quad K_n < 1.338887 (n-1)^{0.526322}.$$

If we use the exact value of  $C_{2^{k_0}}$  instead of estimate (2.1) we can improve (2.2) as  $n$  grows. For example,

$$\begin{aligned} n > 2^6 &\Rightarrow K_n < 1.310883 (n-1)^{0.526322} \\ n > 2^7 &\Rightarrow K_n < 1.306156 (n-1)^{0.526322} \\ n > 2^8 &\Rightarrow K_n < 1.303787 (n-1)^{0.526322}. \end{aligned}$$

**2.2. Complex case.** Let  $(C_n)_{n=1}^\infty$  denote the sequence in [1, Theorem 2.3]. If we fix any  $k_0$ , and as the authors did in [2], we can show that

$$J_n = \begin{cases} C_n & \text{if } n \leq 2^{k_0}, \\ DJ_{\frac{n}{2}} & \text{if } n > 2^{k_0} \text{ is even, and} \\ D \left(J_{\frac{n-1}{2}}\right)^{\frac{n-1}{2n}} \left(J_{\frac{n+1}{2}}\right)^{\frac{n+1}{2n}} & \text{if } n > 2^{k_0} \text{ is odd,} \end{cases}$$

with  $D = e^{\frac{1}{2}-\frac{1}{2}\gamma}$ , satisfies the multilinear Bohnenblust–Hille inequality. For  $n > 2^{k_0}$ , by mimicking the real case we obtain

$$K_n \leq \frac{C_{2^{k_0}}}{D^{k_0-1}} (n-1)^{\log_2 \left(e^{\frac{1}{2}-\frac{1}{2}\gamma}\right)}.$$

Thus, using the values of  $C_{2^k}$  from [1] we have

$$\begin{aligned} n > 2^3 &\Rightarrow K_n < 1.02960973695 (n-1)^{0.304975}, \\ n > 2^4 &\Rightarrow K_n < 1.01089344604 (n-1)^{0.304975}, \\ n > 2^5 &\Rightarrow K_n < 1.00123230777 (n-1)^{0.304975}, \\ n > 2^6 &\Rightarrow K_n < 0.99632125476 (n-1)^{0.304975}, \dots \\ n > 2^{14} &\Rightarrow K_n < 0.99137409768 (n-1)^{0.304975}, \\ n > 2^{15} &\Rightarrow K_n < 0.99136434217 (n-1)^{0.304975}, \dots \\ n > 2^{25} &\Rightarrow K_n < 0.99135459597 (n-1)^{0.304975}, \dots \\ n > 2^{50} &\Rightarrow K_n < 0.99135458644 (n-1)^{0.304975}. \end{aligned}$$

## REFERENCES

- [1] D. Nuñez-Alarcón, D. Pellegrino, J.B. Seoane-Sepúlveda, On the Bohnenblust–Hille inequality and a variant of Littlewoods 4/3 inequality, *J. Funct. Anal.* **264** (2013), 326–336.
- [2] D. Nuñez-Alarcón, D. Pellegrino, J.B. Seoane-Sepúlveda, D.M. Serrano-Rodríguez, There exist multilinear Bohnenblust–Hille constants  $(C_n)_{n=1}^{\infty}$  with  $\lim_{n \rightarrow \infty} (C_{n+1} - C_n) = 0$ , *J. Funct. Anal.* **264** (2013), 429–463.
- [3] D.M. Serrano-Rodríguez, Improving the closed formula for subpolynomial constants in the multilinear Bohnenblust–Hille inequalities, *Linear Alg. Appl.*, in press.

DEPARTAMENTO DE MATEMÁTICA,  
 UNIVERSIDADE FEDERAL DA PARAÍBA,  
 58.051-900 - JOÃO PESSOA, BRAZIL.  
*E-mail address:* danielnunezal@gmail.com

DEPARTAMENTO DE MATEMÁTICA,  
 UNIVERSIDADE FEDERAL DA PARAÍBA,  
 58.051-900 - JOÃO PESSOA, BRAZIL.  
*E-mail address:* dmpellegrino@gmail.com and pellegrino@pq.cnpq.br

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO,  
 FACULTAD DE CIENCIAS MATEMÁTICAS,  
 PLAZA DE CIENCIAS 3,  
 UNIVERSIDAD COMPLUTENSE DE MADRID,  
 MADRID, 28040, SPAIN.  
*E-mail address:* jseoane@mat.ucm.es

DEPARTAMENTO DE MATEMÁTICA,  
 UNIVERSIDADE FEDERAL DA PARAÍBA,  
 58.051-900 - JOÃO PESSOA, BRAZIL.  
*E-mail address:* dmserrano0@gmail.com